
WHAT IS PROLAPSE OF RECTUM?



Rectal prolapsed is when part of the rectum protrudes from the anus. Prolapse occurs when the rectum becomes unattached inside the body and comes out through the anus, effectively turning itself inside out.

TYPES OF PELVIC ORGAN PROLAPSE INCLUDE THE FOLLOWING:

- Cystocele (bladder prolapsed)
- Enterocele (small bowel prolapsed)
- Rectocele (rectum prolapsed)
- Uterine prolapse (prolapse of the uterus)
- Vaginal vault prolapse.

SYMPTOMS OF PROLAPSED

- Sensation of heaviness or pulling in your pelvis.
- Tissue protruding from your vagina.
- Urinary problems, such as urine leakage (incontinence) or urine retention.
- Trouble having a bowel movement.

CAUSES OF WEAKENED PELVIC MUSCLES AND TISSUES INCLUDE:

- Pregnancy.
- Difficult labor and delivery or trauma during childbirth.
- Delivery of a large baby.
- Being overweight or obese.
- Lower estrogen level after menopause.
- Chronic constipation or straining with bowel movements.
- Chronic cough or bronchitis.
- Repeated heavy lifting.

NEUROTHERAPY TREATMENT

Acid treatment formula

(3) Piles point