

## WHAT IS POLIOMYELITIS

Polio, or poliomyelitis, is a crippling and potentially deadly infectious disease. It is caused by the poliovirus. The virus spreads from person to person and can invade an infected person's brain and spinal cord, causing paralysis (can't move parts of the body).

## SYMPTOMS OF POLIO

- Fever.
- Sore throat.
- Headache.
- Vomiting.
- Fatigue.
- Back pain or stiffness.
- Neck pain or stiffness.
- Pain or stiffness in the arms or legs.

## CAUSES OF POLIO

The polio virus usually enters the environment in the feces of someone who is infected. In areas with poor sanitation, the virus easily spreads from feces into the water supply, or, by touch, into food. In addition, because polio is so contagious, direct contact with a person infected with the virus can cause polio.

## NEUROTHERAPY TREATMENT

**First treatment**                      Inflammation treatment formula

Note: If poliomyelitis develops into paralysis, give paralysis treatment

### Second treatment

For the first 3 to 4 months:

First day	I	(30)	Medulla	
				After ½ hour
	II	(6)	Adr	
		(20)	Round arrow	
				After ½ hour
	III	(6)	Adr	
		(30)	Blood supply to legs	

This treatment is to be given on alternate days for 3 to 4 months

Second part    after completing first part for 3-4 months, following sequence should be given

First day A-Heparin, X-Heparin or P-Heparin  
~~III~~

Third day I (3) Gal  
 (7) Liv  
 After ½ hour

Fifth day II 1,25 DCC  
 Normal - Ajay Normal formula

Seventh day Repeat ~~III~~ ~~II~~

Ninth day onwards: ~~III~~

~~III~~

Later give the following treatment:

- (5) Pan
- (3) Wd
- (8) Ch only
- (20) Round Arrow

**Third Treatment**

First day Normal treatment formula  
 Second day Old Genes formula  
 Third day Normal treatment formula

Fourth treatment (3) Pit  
 (6) Adr  
 (3) Swt  
 (6) L5-~~5~~  
 (30) Blood supply to legs